

5.1 – Graphing Exponential Functions

Pg 340
 Math Lab

Exponential Function – A function of the form $y = c(a)^x$, where $c \neq 0, a > 0$

Example 1: a) Fill in the table of values, sketch each graph, and determine the domain and range.

Increasing

a) $y = 10^x$

-3	$1/10^3$ 0.001
-2	$1/10^2$ 0.01
-1	$1/10$ 0.1
0	1
1	10
2	100
3	1000

Increasing

b) $y = 2(5)^x$

-3	0.016
-2	0.08
-1	0.4
0	2
1	10
2	50
3	250

Decreasing





c) $y = (\frac{1}{2})^x$

-3	8
-2	4
-1	2
0	1
1	0.5
2	0.25
3	0.125

Decreasing

d) $y = 8(\frac{1}{4})^x$

-3	512
-2	128
-1	32
0	8
1	2
2	0.5
3	0.125



D: $\{x | x \in \mathbb{R}\}$

R: $\{y | y > 0, y \in \mathbb{R}\}$

Example 2: Determine, from the following table of values, that the function is exponential:

x	y
-3	0.625
-2	1.25
-1	2.5
0	5
1	10
2	20
3	40

$\frac{1.25}{0.625} = 2$
 $\frac{2.5}{1.25} = 2$
 $\frac{5}{2.5} = 2$

$\frac{10}{5} = 2$
 $\frac{20}{10} = 2$
 $\frac{40}{20} = 2$


$y = 5(2)^x$

y-int →

Example 3: Graph $y = (\frac{1}{3})^x$, determine the following: the effect on y when x increases by 1, increasing/decreasing, the intercepts, the equation of any asymptotes, domain, and range.

Decreasing $0 < \frac{1}{3} < 1$
 x-int = none
 y-int = 1
 asymptote: $y = 0$
 D: $\{x | x \in \mathbb{R}\}$
 R: $\{y | y > 0, y \in \mathbb{R}\}$

x	y
-2	9
-1	3
0	1
1	1/3
2	1/9



5.2 – Analyzing Exponential Functions

Pg 349 #3-11

Transformation of Exponential Functions – $y - k = ca^{d(x-h)}$, $a > 0, c \neq 0, d \neq 0$

$|c|$ – Vertical stretch or compression

$\frac{1}{|d|}$ – Horizontal stretch or compression

$c < 0$ – Reflected in the x-axis (flipped vertically)

$d < 0$ – Reflected in the y-axis (flipped horizontally)

k – Vertical translation

h – Horizontal Translation

Exponential Function – The general translation of (x, y) corresponds to $(\frac{x}{d} + h, cy + k)$

Graphing Exponential Functions with Transformations

- 1) Determine lattice points for the original function
- 2) Apply the transformations to your x and y values
- 3) Plot the new points

Example 1: a) Sketch the graph of $y = 2^x$. Determine whether the function is increasing or decreasing, the intercepts, equation of any asymptotes, domain and range.

base = 2

increase: $2 > 1$

x-int = none


y-int = 1

asymptote: $y = 0$

D: $\{x | x \in \mathbb{R}\}$

R: $\{y | y > 0, y \in \mathbb{R}\}$

x	y
-2	1/4
-1	1/2
0	1
1	2
2	4



b) Use the graph of $y = 2^x$ to sketch the graph of $y = 3(2^{-x+2})$. Determine whether the function is increasing or decreasing, the intercepts, equation of any asymptotes, domain and range.

$y = 2^x$

$y = 3(2^{-1(x-2)})$

x	y
-2	0.25
-1	0.5
0	1
1	2
2	4

$y = 0$

-x+2	3y
4	0.75
3	1.5
2	3
1	6
0	12

$y = 0$

d = -1 c = 3
 h = 2 k = 0

Decreasing


x-int = none

y-int = 12

asymptote: $y = 0$

D: $\{x | x \in \mathbb{R}\}$

R: $\{y | y > 0, y \in \mathbb{R}\}$



Natural Logarithm – A logarithm to the base of e where $y = \log_e(x)$ or $y = \ln(x)$

$$\ln(a)^x = x(\ln a)$$

or $\log a^x = x \log a$

Solving for x in Exponential Functions

- 1) Substitute your known value in for y
- 2) Isolate your base and exponent by dividing both sides by a
- 3) Take the natural log (ln) of both sides
- 4) Isolate your x value by arranging knowing that $\ln(a)^x = x(\ln a)$

Example 2: The temperature of cooling water in a cup can be modelled by the equation: $t = 89.726(0.972)^m$, where m is the number of minutes that have passed and t is the temperature in °C. Using this determine:

a) The temperature after 52 minutes

$$t = 89.726(0.972)^{52}$$

$$t = 20.49...$$

b) The time when the water was 51°

$$\frac{51}{89.726} = \frac{89.726(0.972)^x}{89.726}$$

$$\log \frac{51}{89.726} = \log 0.972^x$$

$$\frac{\log \left(\frac{51}{89.726} \right)}{\log(0.972)} = \frac{x \log 0.972}{\log(0.972)}$$

$$19.82... = x$$

Example 3: For every metre below the surface of water, the light intensity is reduced by 2.5%. The percent, P , of light remaining at a depth d metres can be modelled by the function: $P = 100(0.975)^d$

a) Sketch the graph of the function, determine whether the function is increasing or decreasing, the intercepts, equation of any asymptotes, domain and range.

decrease $0 < 0.9 < 1$

asymptote: $y = 0$

x-int = none

domain: $\{x \mid x \in \mathbb{R}\}$

y-int = 100

range: $\{y \mid y > 0, y \in \mathbb{R}\}$

b) To the nearest percent, how much light remains at a depth of 10 m?

$$P = 100(0.975)^{10}$$

$$P = 77.63...$$

c) To the nearest metre, what is the depth when only 50% of the light remains?

$$\frac{50}{100} = \frac{100(0.975)^x}{100}$$

$$\log 0.5 = \log 0.975^x$$

$$\frac{\log 0.5}{\log 0.975} = \frac{x \log 0.975}{\log 0.975}$$

$$x = 27.4...$$

5.3 – Solving Exponential Equations Page 364 # 3-14

- Product of Powers** – Multiplying variables with exponents: $x^a \cdot x^b = x^{a+b}$ $2^7 \cdot 2^3 = 2^{10}$
- Quotient of Powers** – Dividing variables with exponents: $\frac{x^a}{x^b} = x^{a-b}$ $\frac{2^2}{2^1} = 2^1$
- Power of a Power** – Variable with an exponent to the power of an exponent: $(x^a)^b = x^{a \cdot b}$ $(2^3)^3 = 2^9$
- Power of a Quotient** – A fraction raised to an exponent: $\left(\frac{x}{y}\right)^a = \frac{x^a}{y^a}$ $\left(\frac{2}{3}\right)^2 = \frac{2^2}{3^2}$
- Negative Exponent** – An exponent that is negative becomes positive with the reciprocal of the base: $x^{-a} = \frac{1}{x^a}$ $\frac{1}{2^2} = 2^{-2}$
- Fractional Exponent** – A fractional exponent can be written as a radical: $x^{\frac{m}{n}} = \sqrt[n]{x^m}$ $2^{\frac{3}{2}} = \sqrt{2^3}$

Exponential Equation – Contains a power with a variable in the exponent.

Solving Exponential Equations Using Common Bases

- 1) Determine the common base on the right side and left side of the equation
- 2) Divide your number by the common base, counting how many times you've done this, until you get to 1
- 3) Rewrite this number as a common base to the power of the amount of times you were able to divide
- 4) Make sure the common bases are the same, then you can cancel them out, leaving only the exponents
- 5) Solve for x

Example 1: Solve each equation:

a) $4^x = \frac{1}{256}$

$256 \div 4 = 4 = 4 \div 4 = 1$

$4^4 = 256$

$4^x = \frac{1}{4^4}$

$4^x = 4^{-4}$

$x = -4$

b) $27^x = 9^{2x-1}$

$3^{3(x)} = 3^{2(2x-1)}$

$3x = 2(2x-1)$

$3x^{+2} = 4x - 2^{+2}$

$-3x \quad -3x$

$2 = x$

If you want another example of this go to Example 1 on page 359.

Example 2: Solve each equation:

a) $2^x = 8\sqrt[3]{2}$

$2^x = 2^3 \cdot 2^{\frac{1}{3}}$

$2^x = 2^{3+\frac{1}{3}}$

$2^x = 2^{10/3}$

$x = \frac{10}{3}$

b) $(\sqrt{125})^{2x+1} = \sqrt[3]{625}$

$(\sqrt{5^3})^{2x+1} = \sqrt[3]{5^4}$

$5^{\frac{3}{2}(2x+1)} = 5^{4/3}$

$\left(\frac{3}{2}(2x+1)\right) = \left(\frac{4}{3}\right)$

$9(2x+1) = 8$

$18x + 9 = 8^{-9}$

$\frac{18x}{18} = \frac{-1}{18}$

$x = -\frac{1}{18}$

If you want another example of this go to Example 2 on page 360.

Compound Interest – The interest that is earned or paid on both the principal and the accumulated interest.

Compounding Period – The time over which interest is determined

Number of Compounding Periods in a Year					
Annually	Semi-annually	Quarterly	Monthly	Weekly	Daily
1	2	4	12	52	365

Principal – The initial amount that money is invested or loaned.

Exponential Growth – A function that models exponential growth has the form of $y = ak^{bx}$, where $k^b > 1$, and $a \in R, b \in R, k > 0$. K is the growth factor.

Compound Interest – $A = P \left(1 + \frac{i}{n}\right)^{nt}$, where A is future value, P is the principle (amount invested, or A_0), i is interest rate, n is compounding periods and t is the term.

Example 3: A principal of \$1500 is invested at 4% annual interest, compounded quarterly. To the nearest quarter of a year, when will the amount be \$2500?

$$\begin{aligned}
 A &= 2500 \\
 P &= 1500 \\
 i &= 0.04 \\
 n &= 4 \\
 t &= ?
 \end{aligned}$$

$$A = P \left(1 + \frac{i}{n}\right)^{nt}$$

$$\frac{2500}{1500} = \frac{1500 \left(1 + \frac{0.04}{4}\right)^{4t}}{1500}$$

$$\log\left(\frac{2500}{1500}\right) = \log\left(1 + \frac{0.04}{4}\right)^{4t}$$

$$\frac{\log\left(\frac{2500}{1500}\right)}{4 \log\left(1 + \frac{0.04}{4}\right)} = \frac{4t \log\left(1 + \frac{0.04}{4}\right)}{4 \log\left(1 + \frac{0.04}{4}\right)}$$

$$t = 12.83 \text{ years}$$

If you want another example of this go to Example 3 on page 362.

Exponential Decay – A function that models exponential decay has the form of $y = ak^{bx}$, where $0 < k^b < 1$, and $a \in R, b \in R, k > 0$. K is the decay factor.

Example 4: The function $P = 101.3(0.88)^h$ models the atmospheric pressure, P kilopascals, at an altitude of h kilometres. If the cabin pressure in an airplane is less than 70 kPa, passengers can suffer from altitude sickness. To the nearest kilometre, at what altitude is the atmospheric pressure 70 kPa?

$$\begin{aligned}
 P &= 101.3(0.88)^h \\
 70 &= \frac{101.3(0.88)^h}{101.3} \\
 \log\left(\frac{70}{101.3}\right) &= \log(0.88)^h \\
 \frac{\log\left(\frac{70}{101.3}\right)}{\log(0.88)} &= \frac{h \log(0.88)}{\log(0.88)} \\
 2.891\dots &= h
 \end{aligned}$$

If you want another example of this go to Example 4 on page 363.


5.4 – Logarithms and the Logarithmic Function

Page 381
 # 4-8, 11, 13, 15, 16

Example 1: Graph the functions $y = 10^x$ and $y = \log_{10}x$. Determine whether the function is increasing or decreasing, the intercepts, equation of any asymptotes, domain and range of each function. How are these graphs related?

-2	1/100
-1	1/10
0	1
1	10
2	100

-1	und
0	und
0.01	-2
0.1	-1
1	0
10	1
100	2



increasing
 x-int = none
 y-int = 1
 asymptote: $y = 0$
 D: $\{x \mid x \in \mathbb{R}\}$
 R: $\{y \mid y > 0, y \in \mathbb{R}\}$

increasing
 x-int = 1
 y-int = none
 asymptote: $x = 0$
 D: $\{x \mid x > 0, x \in \mathbb{R}\}$
 R: $\{y \mid y \in \mathbb{R}\}$

Logarithmic Function – The logarithm of a number is an exponent. $\log_b c = a$ is the power to which b is raised to get c . The base of the logarithm is the same as the base of the power. When $\log_b c = a$, then $c = b^a$, where $b > 0, b \neq 1, c > 0$.

Common Logarithm – Numbers based on powers of 10, $\log_{10}x$ is called the common logarithm. When logarithms to base 10 are written, the base is often not shown; so $\log_{10}x$ is written as $\log x$.

$$y = \log_b x \text{ is equivalent to } x = b^y$$

Changing Form Between Exponential and Logarithmic

- 1) Determine the base, rewrite as the new base
- 2) Invert the x and y values

Example 2: Write each exponential expression as a logarithmic expression:

a) $3^3 = 27$

$$\log_3 27 = 3$$

b) $5^{-2} = \frac{1}{25}$

$$\log_5 \frac{1}{25} = -2$$

c) $4^0 = 1$

$$\log_4 1 = 0$$

Write each logarithmic expression as an exponential expression:

a) $\log_3 81 = 4$

$$3^4 = 81$$

b) $\log_5 125 = 3$

$$5^3 = 125$$

c) $\log_6 1 = 0$

$$6^0 = 1$$

Evaluating Logarithms

- 1) Rewrite as an exponential equation
- 2) Simplify

Example 3: Evaluate each logarithm:

a) $\log_5 3125 \rightarrow$ Change to base 5
 $3125 \div 5 \div 5 \div 5 \div 5 \div 5 = 1$
 $\log_5 5^5 = x$
 $5^x = 5^5$
 $x = 5$

b) $\log_6 \left(\frac{1}{216}\right)$
 $\log_6 \left(\frac{1}{6^3}\right)$
 $\log_6 (6^{-3}) = x$
 $6^x = 6^{-3}$
 $x = -3$

c) $\log_8 (2\sqrt[3]{2})$
 $\log_{2^3} (2 \cdot 2^{1/3})$
 $\log_{2^3} (2^{1+1/3})$
 $\log_{2^3} (2^{4/3}) = x$
 $2^{3(x)} = 2^{4/3}$
 $\frac{3x}{3} = \frac{4/3}{3}$

If you want another example of this go to Example 2 on page 378.

Example 4: To the nearest tenth, estimate the value of $\log_5 100$ to 2 decimal places.

$5^x = 100$
 try 2.8
 $5^{2.8} = 90.597$
 $5^{2.9} = 106.417$
 $5^2 = 25$
 $5^3 = 125$
 ← closer to this
 $5^{2.85} = 98.189$
 $5^{2.86} = 99.782$
 $5^{2.87} = 101.401$
 $x = 4/9$
 $\log_5 100 \approx 2.86$

If you want another example of this go to Example 3 on page 379.

Sketching Graphs of Logarithmic Functions

- 1) Determine the inverse Exponential Function
- 2) Create a table of values for the exponential function
- 3) Invert your x and y values to sketch the points

Example 5: Graph $y = \log_4 x$. Identify the intercepts, the equations of any asymptotes, and the domain and range of the function.


x	y
-2	1/16
-1	1/4
0	1
1	4
2	16

$y = 0$

x	y
1/16	-2
1/4	-1
1	0
4	1
16	2

$x = 0$

$y = c \log_a x$
 x-int (1, 0)
 (a, c)



If you want another example of this go to Example 4 on page 379.

Page 393
 # 4-8, 11, 12, 13, 16

5.5 – The Laws of Logarithm

- Product Law:** $\log_b xy = \log_b x + \log_b y$, $b > 0, b \neq 1$
Quotient Law: $\log_b \left(\frac{x}{y}\right) = \log_b x - \log_b y$, $b > 0, b \neq 1$
Power Law: $\log_b x^k = k \log_b x$, $b > 0, b \neq 1, k \in R$

Example 1: Use a law of logarithms to simplify each expression. Use a calculator to verify the answers:

- | | | |
|----------------------|---------------|----------------------------------|
| a) $\log 7 + \log 8$ | b) $5 \log 2$ | c) $\log 80 - \log 16$ |
| $\log(7 \cdot 8)$ | $\log 2^5$ | $\log\left(\frac{80}{16}\right)$ |
| $\log(56)$ | $\log(32)$ | $\log 5$ |
| 1.748... | 1.505... | 0.698... |

If you want another example of this go to Example 1 on page 390.

Example 2: Write each expression as a single logarithm

- | | | |
|------------------------|-------------------------------------|----------------------------------|
| a) $\log x + 3 \log y$ | b) $\log x + 2 \log y - 4 \log z$ | c) $\log_2 6 - 3$ |
| $\log x + \log y^3$ | $\log(x) + \log(y^2) - \log(z^4)$ | $\log_2 6 - \log_2 2^3$ |
| $\log(x \cdot y^3)$ | $\log\left(\frac{xy^2}{z^4}\right)$ | $\log_2 6 - \log_2 8$ |
| | | $\log_2\left(\frac{6}{8}\right)$ |

If you want another example of this go to Example 2 on page 390.

Example 3: Write each expression in terms of $\log a$, $\log b$, and/or $\log c$.

- | | |
|-------------------------------------|-----------------------------------------------------|
| a) $\log\left(\frac{a}{b^2}\right)$ | b) $\log\left(\frac{a^2 b^{\frac{1}{3}}}{c}\right)$ |
| $\log a - \log b^2$ | $\log a^2 + \log b^{\frac{1}{3}} - \log c$ |
| $\log a - 2 \log b$ | $2 \log a + \log \sqrt[3]{b} - \log c$ |

If you want another example of this go to Example 3 on page 391.

Example 4: Evaluate each expression:

- | | |
|-------------------------------------|------------------------------------------|
| a) $3 \log_9 6 - \log_9 72$ | b) $2 \log_4 6 - 3 \log_4 3 + \log_4 12$ |
| $\log_9 6^3 - \log_9 72$ | $\log_4 6^2 - \log_4 3^3 + \log_4 12$ |
| $\log_9 216 - \log_9 72$ | $\log_4 36 - \log_4 27 + \log_4 12$ |
| $\log_9\left(\frac{216}{72}\right)$ | $\log_4\left(\frac{36}{27}\right)(12)$ |
| $\log_9 3$ | $\log_4(16)$ |
| $9^x = 3$ | $4^x = 16$ |
| $3^{2x} = 3^1$ | |
| $\frac{2x}{2} = \frac{1}{2}$ | |
| $x = \frac{1}{2}$ | $4^x = 4^2$ |
| | $x = 2$ |

If you want another example of this go to Example 4 on page 392.

5.6 – Analyzing Logarithmic Functions Page 405

3 - 12

Example 1: Solve for y in $y = \log_b x$

$$\begin{aligned}
 b^y &= x \\
 \log b^y &= \log x \\
 y \log b &= \frac{\log x}{\log b} \\
 y &= \frac{\log x}{\log b}
 \end{aligned}$$

Changing Base of a Logarithm – $\log_b x = \frac{\log x}{\log b}$, where $a, b > 0, b \neq 1; x > 0$

Example 2: Approximate the value of each logarithm, to the nearest thousandth. Write the related exponential expression.

a) $\log_5 50$

$$\frac{\log 50}{\log 5} \approx 2.430\dots$$

b) $\log_8 6$

$$\frac{\log 6}{\log 8} \approx 0.8616\dots$$

If you want another example of this go to Example 1 on page 402.

Example 3: Fill in the table of values, sketch each graph, and determine the domain and range.

increasing

$y = \log_{10} x$	
-1	und
0	und
0.5	-
1	0
10	1

increasing





$y = 2\log_{10} x$	
-1	und
0	und
0.5	-
1	0
10	2

decreasing

$y = -3\log_{10} x$	
-1	und
0	und
0.5	+
1	0
10	-3

decreasing

$y = \log_8 x$	
-1	und
0	und
0.5	-
1	0
8	1



$D: \{x \mid x > 0, x \in \mathbb{R}\}$ $R: \{y \mid y \in \mathbb{R}\}$

Transformation of Logarithmic Functions – $y - k = c \log_a d(x - h)$, $a > 0, c \neq 0, d \neq 0$

- $|c|$ – Vertical stretch or compression
- $\frac{1}{|d|}$ – Horizontal stretch or compression
- $c < 0$ – Reflected in the x-axis (flipped vertically)
- $d < 0$ – Reflected in the y-axis (flipped horizontally)
- k – Vertical translation
- h – Horizontal Translation

Logarithmic Function – The general translation of (x, y) corresponds to $(\frac{x}{d} + h, cy + k)$

Sketching Graphs of Logarithmic Functions

- 1) Use lattice points for the x-intercept and (a,1)
- 2) Determine any transformations made
- 3) Transform the points

$$y - K = c \log_a d(x - h)$$

$$\left(\frac{x}{d} + h, cy + K\right)$$

Example 4: Sketch the graph of $y = \log_3 x$ and $y = \log_3(2x + 6)$

$\log_3 x$	
x	y
1	0
3	1

$x = 0$

$\log_3 2(x+3)$	
$\frac{x}{a} - 3$	y
-2.5	0
-1.5	1

$x = -3$


$d = 2$ $c = 1$
 $h = -3$ $K = 0$

$$y = \log_3 2(0+3)$$

$$y = \log_3 2(3)$$

$$y = \log_3 6$$

$$y = \frac{\log 6}{\log 3} = 1.63$$



Increasing
 x-int = -2.5
 y-int = 1.63...
 asymptote: $x = -3$

$D: \{x \mid x > -3, x \in \mathbb{R}\}$
 $R: \{y \mid y \in \mathbb{R}\}$

Example 5: Sketch the graph of $y = \log_2 x$ and $y = \log_2 2x - 1$

$\log_2 x$	
x	y
1	0
2	1


$x = 0$

$\log_2 2x - 1$	
$\frac{x}{a}$	y - 1
0.5	-1
1	0

$x = 0$

$d = 2$ $c = 1$
 $h = 0$ $K = -1$

increasing
 x-int = 1
 y-int = none
 asymptote: $x = 0$
 $D: \{x \mid x > 0, x \in \mathbb{R}\}$
 $R: \{y \mid y \in \mathbb{R}\}$



Example 6: Sketch the graph of $y = \log_5 x$ and $y = -\frac{1}{2} \log_5(2x + 2)$

$\log_5 x$

x	y
1	0
5	1

$x=0$

$$y = -\frac{1}{2} \log_5 2(x+1)$$

$\frac{x}{2} - 1$	$-\frac{y}{2}$
-0.5	0
1.5	-0.5

$x = -1$


$d = 2$ $c = -1/2$
 $h = -1$ $k = 0$

$$y = -\frac{1}{2} \log_5 2(0+1)$$

$$y = -0.5 \log_5(2)$$

$$y = -0.5 \left(\frac{\log 2}{\log 5} \right)$$

$$y = -0.215\dots$$



decreasing
 $x\text{-int} = -0.5$
 $y\text{-int} = -0.215\dots$
 asymptote: $x = -1$
 $D: \{x \mid x > -1, x \in \mathbb{R}\}$
 $R: \{y \mid y \in \mathbb{R}\}$

Example 7: Sketch the graph of $y = \log_2 x$ and $y = 2 \log_2 \left(-\frac{1}{3}(x+1) + 2\right)$

$\log_2 x$

x	y
1	0
2	1

$x=0$

$$y = 2 \log_2 \left(-\frac{1}{3}(x+1) + 2\right)$$

$-3x - 1$	$2y + 2$
-4	2
-7	4

$x = -1$

$d = -\frac{1}{3}$ $c = 2$
 $h = -1$ $k = 2$

$$-\frac{0}{2} = 2 \log_2 \left(-\frac{1}{3}(x+1) + 2\right) - 2$$


$$-\frac{2}{2} = \frac{2 \log_2 \left(-\frac{1}{3}(x+1) + 2\right)}{2}$$

$$-1 = \log_2 \left(-\frac{1}{3}(x+1) + 2\right)$$

$$2^{-1} = -\frac{1}{3}(x+1) + 2$$

$$-\frac{1}{2} = -\frac{1}{3}(x+1) + 2$$

$$-2.5 = x$$



decreasing
 $x\text{-int} = -2.5$
 $y\text{-int} = \text{none}$
 asymptote: $x = -1$
 $D: \{x \mid x < -1, x \in \mathbb{R}\}$
 $R: \{y \mid y \in \mathbb{R}\}$

5.7 – Solving Logarithmic and Exponential Functions

Logarithmic Equation – An equation that contains the logarithm of a variable. The laws of logarithms may be used to solve logarithmic equations.

Example 1: Solve: $\log_3 9x + \log_3 x = 4$. Verify the solution.

$$\log_3 (9x)x = 4$$

$$\log_3 9x^2 = 4$$

$$3^4 = 9x^2$$

$$\frac{81}{9} = \frac{9x^2}{9}$$

$$\sqrt{9} = \sqrt{x^2}$$

$$\pm 3 = x$$

$$x = 3$$
~~$$x = -3$$~~

extraneous because you can't take the log of a negative number so it becomes undefined at $\log_3 9x$ and $\log_3 x$

If you want another example of this go to Example 1 on page 418.

Example 2: Solve then verify each equation.

a) $\log 6x = \log(x+6) + \log(x-1) - \log 6x$

b) $3 = \log_2(x+2) + \log_2 x$

$$0 = \log(x+6) + \log(x-1) - \log 6x$$

$$0 = \log_{10} \left(\frac{(x+6)(x-1)}{6x} \right)$$

$$10^0 = \frac{(x+6)(x-1)}{6x}$$

$$1 = \frac{x^2 + 5x - 6}{6x}$$

$$6x = x^2 + 5x - 6$$

$$0 = x^2 - x - 6$$

$$0 = (x-3)(x+2)$$

$$x = 3 \quad \del{x = -2}$$

becomes undefined at $\log 6x$ and $\log(x-1)$

$$3 = \log_2(x+2)(x)$$

$$3 = \log_2(x^2 + 2x)$$

$$2^3 = x^2 + 2x$$

$$8 = x^2 + 2x - 8$$

$$-8 = x^2 + 2x - 8$$

$$0 = (x+4)(x-2)$$

$$\del{x = -4} \quad x = 2$$

↓
 is undefined at $\log_2(x+2)$
 $\log_2 x$

$$3 = \log_2(x+2)(x)$$

$$\log_2 2^3 = \log_2(x^2 + 2x)$$

$$8 = x^2 + 2x - 8$$

$$-8 = x^2 + 2x - 8$$

$$0 = x^2 + 2x - 8$$

$$(x+4)(x-2)$$

$$x = -4$$

$$x = 2$$

If you want another example of this go to Example 2 on page 419.

Example 3: Solve each exponential equation algebraically. Give the solution to the nearest hundredth.

a) $4^x = 12$

$$\log_4 12 = x$$

* change it to a logarithm

$$\frac{\log 12}{\log 4} = x$$

$$1.792\dots$$

OR

$$\log 4^x = \log 12$$

* take log of both sides

$$\frac{x \log 4}{\log 4} = \frac{\log 12}{\log 4}$$

$$x = \frac{\log 12}{\log 4}$$

$$x = 1.792\dots$$

c) $3^{x+1} = 6^x$

$$\log 3^{x+1} = \log 6^x$$

$$(x+1)\log 3 = x \log 6$$

$$x \log 3 + 1 \log 3 = x \log 6 - x \log 3$$

$$-x \log 3$$

$$\log 3 = x(\log 6 - \log 3)$$

$$\log 3 = x \left(\log \left(\frac{6}{3} \right) \right)$$

$$\frac{\log 3}{\log 2} = \frac{x \log 2}{\log 2}$$

$$1.58\dots = x$$

b) $3(2^{x+1}) = \frac{36}{3}$

$$2^{x+1} = 12$$

$$\log_2 12 = x+1$$

$$\frac{\log 12}{\log 2} - 1 = x+1$$

$$2.5849\dots = x$$

OR

$$\log 2^{x+1} = \log 12$$

$$(x+1) \frac{\log 2}{\log 2} = \frac{\log 12}{\log 2}$$

$$x+1 = \frac{\log 12}{\log 2} - 1$$

$$x = 2.5849\dots$$

d) $2^{x+3} = 6^{x-1}$

$$\log 2^{x+3} = \log 6^{x-1}$$

$$(x+3)\log 2 = (x-1)\log 6$$

$$x \log 2 + 3 \log 2 = x \log 6 - \log 6$$

$$-x \log 2$$

$$3 \log 2 + \log 6 = x \log 6 - x \log 2$$

$$\log 2^3 + \log 6 = x(\log 6 - \log 2)$$

$$\log 8 + \log 6 = x \left(\log \left(\frac{6}{2} \right) \right)$$

$$\log(8 \cdot 6) = x \log 3$$

$$\frac{\log 48}{\log 3} = \frac{x \log 3}{\log 3}$$

$$\frac{\log 48}{\log 3} = x$$

$$3.52\dots = x$$

5.8 – Solving Problems with Exponents and Logarithms

Page 435

3-7, 9, 10

Annuity – A continual payment of a fixed amount. Solved using the formula:

$$\text{Investments: } A = P \left[\frac{\left(1 + \frac{i}{n}\right)^{nt} - 1}{\frac{i}{n}} \right]$$

$$\text{Loans: } A = P \left[\frac{1 - \left(1 + \frac{i}{n}\right)^{-nt}}{\frac{i}{n}} \right]$$

Example 1: Determine how many monthly investments of \$200 would have to be made into an account that pays 6% annual interest, compounded monthly, for the future value to be \$100 000.

$A = 100\,000$
 $P = 200$
 $i = 0.06$
 $n = 12$
 $t = ?$
 $1 + \frac{i}{n} = 1.005$

$$A = P \left(\frac{\left(1 + \frac{i}{n}\right)^{nt} - 1}{\frac{i}{n}} \right)$$

$$\frac{100\,000}{200} = \frac{200}{200} \left(\frac{(1.005)^{12t} - 1}{0.005} \right)$$

$$500 = \frac{(1.005)^{12t} - 1}{0.005} \times 0.005$$

$$2.5 = (1.005)^{12t} - 1$$

$$3.5 = (1.005)^{12t}$$

$$\log 3.5 = \log (1.005)^{12t}$$

$$\frac{\log 3.5}{12 \log 1.005} = \frac{12t \log (1.005)}{12 \log 1.005}$$

$$20.93... = t$$

(years)

$$\times 12 = 251.178 \text{ months}$$

If you want another example of this go to Example 1 on page 431.

Example 2: A person borrows \$15 000 to buy a car. The person can afford to pay \$300 a month. The loan will be repaid with equal monthly payments at 6% annual interest, compounded monthly. How many monthly payments will the person make?

$A = 15\,000$
 $P = 300$
 $i = 0.06$
 $n = 12$
 $t = ?$
 $1 + \frac{i}{n} = 1.005$

$$A = P \left(\frac{1 - \left(1 + \frac{i}{n}\right)^{-nt}}{\frac{i}{n}} \right)$$

$$\frac{15\,000}{300} = \frac{300}{300} \left(\frac{1 - (1.005)^{-12t}}{0.005} \right)$$

$$50 = \frac{1 - (1.005)^{-12t}}{0.005} \times 0.005$$

$$0.25 = \frac{1 - 1.005^{-12t}}{1}$$

$$-0.75 = -1.005^{-12t}$$

$$\log -0.75 = \log 1.005^{-12t}$$

$$\frac{\log 0.75}{-12 \log (1.005)} = \frac{-12t \log (1.005)}{-12 \log (1.005)}$$

$$4.806... \text{ (years)} = t$$

$$\times 12$$

$$57.68... \text{ (monthly payments)}$$

$$\times \$300$$

$$17\,304.04 \text{ total paid}$$

$$- 15\,000$$

$$= \$2\,304.04 \text{ interest paid}$$

If you want another example of this go to Example 2 on page 432.

Richter Scale – The intensity of vibrations of an earthquake, I microns, is measured 100 km away from the epicentre of the earthquake. The intensity is compared to the intensity, S , of a standard earthquake. The logarithmic scale for measuring the intensity is: $M = \log \left(\frac{I}{S} \right)$

Example 3: The most intense earthquake ever recorded was in Chile in May 1960, with a magnitude of 9.5. In January of 2010, Haiti experienced an earthquake with a magnitude of 7.0. Calculate the intensity of the earthquake in Chile and Haiti. How many times as intense as the Haiti earthquake was the Chile earthquake?

Haiti

$$M = \log \frac{I}{S}$$

$$7 = \log \frac{I}{S}$$

$$10^7 \times S = I$$

Chile

$$M = \log \frac{I}{S}$$

$$9.5 = \log \frac{I}{S}$$

$$10^{9.5} = \frac{I}{S} \times S$$

$$10^{9.5} S = I$$

$$\frac{\text{Chile}}{\text{Haiti}} = \frac{10^{9.5} S}{10^7 S}$$

$$= 10^{2.5} \Rightarrow 316.227 \text{ times as intense as Haiti}$$

If you want another example of this go to Example 2 on page 432.